

**Geometry**  
**Second Semester 2010/11**  
Homework 7

1. Find the velocity, acceleration and speed of each of the following parametrised curves  $\alpha: \mathbf{R} \rightarrow \mathbf{R}^{n+1}$ .
  - (a)  $n = 1$  and  $\alpha(t) = (t, t^2)$ ,
  - (b)  $n = 1$  and  $\alpha(t) = (\cos t, \sin t)$ ,
  - (c)  $n = 1$  and  $\alpha(t) = (\cos 3t, \sin 3t)$ ,
  - (d)  $n = 2$  and  $\alpha(t) = (\cos t, \sin t, t)$ , and
  - (e)  $n = 3$  and  $\alpha(t) = (\cos t, \sin t, 2 \cos t, 2 \sin t)$ .
2. Show that if  $\alpha: I \rightarrow \mathbf{R}^{n+1}$  is a parametrised curve with constant speed, then  $\ddot{\alpha}(t) \perp \dot{\alpha}(t)$  for all  $t \in I$ .
3. Let  $\alpha: I \rightarrow \mathbf{R}^{n+1}$  be a parametrised curve such that  $\dot{\alpha}(t) \neq \mathbf{0}$  for all  $t \in I$ . Show that there exists a unit speed reparametrisation  $\beta$  of  $\alpha$ . That is, show that there exists an interval  $J$  and a smooth surjective function  $h: J \rightarrow I$  such that  $h' > 0$  and  $\beta = \alpha \circ h$  has unit speed. [Hint: Set  $h = s^{-1}$ , where  $s(t) = \int_{t_0}^t \|\alpha(\tau)\| d\tau$  for some  $t_0 \in I$ .]
4. Let  $\mathbf{X}$  and  $\mathbf{Y}$  be smooth vector fields along the parametrised curve  $\alpha: I \rightarrow \mathbf{R}^{n+1}$  and let  $f: I \rightarrow \mathbf{R}$  be a smooth function along  $\alpha$ . Verify the following identities, which we claimed to hold in class:
  - (a)  $(\mathbf{X} + \mathbf{Y})' = \dot{\mathbf{X}} + \dot{\mathbf{Y}}$ ;
  - (b)  $(f\mathbf{X})' = f'\mathbf{X} + f\dot{\mathbf{X}}$ ; and
  - (c)  $(\mathbf{X} \cdot \mathbf{Y})' = \dot{\mathbf{X}} \cdot \mathbf{Y} + \mathbf{X} \cdot \dot{\mathbf{Y}}$ .
5. Let  $S$  be an  $n$ -surface in  $\mathbf{R}^{n+1}$ ,  $p \in S$ ,  $\mathbf{v} \in S_p$  and let  $\alpha: I \rightarrow S$  be the maximal geodesic in  $S$  passing through  $p$  with velocity  $\mathbf{v}$ . Show that the maximal geodesic  $\beta$  in  $S$  with  $\beta(0) = p$  and  $\dot{\beta}(0) = c\mathbf{v}$ , with  $c \in \mathbf{R}$ , is given by the formula  $\beta(t) = \alpha(ct)$  for all  $t \in \tilde{I}$ .
6. Let  $\alpha: I \rightarrow S$  be a geodesic on an  $n$ -surface  $S$  and let  $\beta = \alpha \circ h$ , where  $h: \tilde{I} \rightarrow I$  is a smooth surjective function, be a reparametrisation of  $\alpha$ . Show that  $\beta$  is a geodesic on  $S$  if and only if  $h(t) = at + b$  for some  $a, b \in \mathbf{R}$  and all  $t \in \tilde{I}$ .
7. An  $n$ -surface in  $\mathbf{R}^{n+1}$  is said to be *geodesically complete* if every maximal geodesic on  $S$  has domain  $\mathbf{R}$ . Which of the following  $n$ -surfaces are geodesically complete?
  - (a) The  $n$ -sphere  $\mathbf{S}^n \subset \mathbf{R}^{n+1}$  (see Example 4.2).
  - (b)  $f^{-1}(1) \subset \mathbf{R}^{n+1}$  where  $f: U \rightarrow \mathbf{R}$  has domain  $U = \{(x_1, x_2, \dots, x_{n+1}) \mid x_{n+1} < 1\}$  and is given by  $f(x_1, x_2, \dots, x_{n+1}) = x_1^2 + x_2^2 + \dots + x_{n+1}^2$  for each  $(x_1, x_2, \dots, x_{n+1}) \in U$ . This is the  $n$ -sphere  $\mathbf{S}^n \setminus \{(0, \dots, 0, 1)\}$  with the north pole deleted.
  - (c)  $f^{-1}(0) \subset \mathbf{R}^3$  where  $f: U \rightarrow \mathbf{R}$  has domain  $U = \{(x_1, x_2, x_3) \mid x_3 > 1\}$  and is given by  $f(x_1, x_2, x_3) = x_1^2 + x_2^2 - x_3^2$  for each  $(x_1, x_2, \dots, x_{n+1}) \in U$ . This is a cone with the vertex deleted.
  - (d) The 2-surface  $S$  in  $\mathbf{R}^3$  which is the cylinder over the plane curve  $\mathbf{S}^1$  (see Examples 4.2 and 4.5).