

Partial Differential Equations (TATA27)
Spring Semester 2019
Homework 2

Preparation for the next seminar

In preparation for Seminar 3 read through Chapter 4 and Section 5.1 and attempt the following two problems.

- 2.1 Let $\mathbf{R}_+^2 = \mathbf{R} \times (0, \infty)$, $C(\overline{\mathbf{R}_+^2})$ denote the set of continuous real-valued functions on $\overline{\mathbf{R}_+^2}$ and $C^1(\mathbf{R}_+^2)$ denote the set of continuously differentiable real-valued functions on \mathbf{R}_+^2 . Consider the boundary-value problem

$$\begin{cases} u_x(x, y) + yu_y(x, y) = 0 & \text{for all } (x, y) \in \mathbf{R}_+^2, \text{ and} \\ u(x, 0) = \phi(x) & \text{for all } x \in \mathbf{R}. \end{cases}$$

- (a) Show that if $\phi(x) = x$ for all $x \in \mathbf{R}$, then no solution exists in $C(\overline{\mathbf{R}_+^2}) \cap C^1(\mathbf{R}_+^2)$.
(b) Show that if $\phi(x) = 1$ for all $x \in \mathbf{R}$, then there are many solutions in $C(\overline{\mathbf{R}_+^2}) \cap C^1(\mathbf{R}_+^2)$.
- 2.2 Fix $\ell > 0$ and consider the following boundary-value problem. Given a function $f: (0, \ell) \rightarrow \mathbf{R}$ we wish to find $u: [0, \ell] \rightarrow \mathbf{R}$ which is twice continuously differentiable such that

$$\begin{cases} u''(x) + u'(x) = f(x) & \text{for all } x \in (0, \ell), \text{ and} \\ u'(0) = u(0) = \frac{1}{2}(u'(\ell) + u(\ell)). \end{cases}$$

- (a) Prove that if a solution u exists, it is not unique.
(b) Find two conditions we must place on f for a solution to exist.

Group work

We will work on the following exercise at the end of the seminar then we will discuss possible solutions together in Seminar 4.

- 2.3 Let Ω be a bounded open set. Prove that continuous functions $u: \overline{\Omega} \rightarrow \mathbf{R}$ which satisfy

$$\Delta u(\mathbf{x}) + \mathbf{x} \cdot \nabla u(\mathbf{x}) \geq 0$$

for $\mathbf{x} \in \Omega$ also satisfy the weak maximum principle:

$$\max_{\overline{\Omega}} u = \max_{\partial\Omega} u.$$

Review exercises

Here's a couple of additional exercises for you to try.

- 2.4 Suppose $u: \mathbf{R}^2 \rightarrow \mathbf{R}$ is a harmonic function.

- (a) For constants $a, b \in \mathbf{R}$ show that $v: \mathbf{R}^2 \rightarrow \mathbf{R}$ defined by

$$v(x, y) = u(x + a, y + b) \quad \text{for all } x, y \in \mathbf{R}$$

is harmonic.

- (b) For a constant $\alpha \in \mathbf{R}$ show that $w: \mathbf{R}^2 \rightarrow \mathbf{R}$ defined by

$$w(x, y) = u(x \cos \alpha + y \sin \alpha, y \cos \alpha - x \sin \alpha) \quad \text{for all } x, y \in \mathbf{R}$$

is harmonic.

This exercise shows that Laplace's equation in the plane is invariant under rigid motions (translations and rotations).

2.5 The Schrödinger equation is a good model for the behaviour of particles at the atomic and subatomic level. Solutions $u: \mathbf{R}^3 \times \mathbf{R} \rightarrow \mathbf{C}$ are complex-valued and are related to the probability that a particle can be found in a specific region. The equation which models the motion of an electron around a hydrogen nucleus has the form

$$-i\hbar \frac{\partial u}{\partial t}(\mathbf{x}, t) = \frac{\hbar^2}{2m} \Delta u(\mathbf{x}, t) + \frac{e^2}{|\mathbf{x}|} u(\mathbf{x}, t)$$

for real constants \hbar , m and e and all $\mathbf{x} \in \mathbf{R}^3$ and $t \in \mathbf{R}$. Assume that u and $\partial_t u$ are continuous functions, and u , $\partial_t u$ and ∇u satisfy the estimate $|u(\mathbf{x}, t)|^2 + |\partial_t u(\mathbf{x}, t)|^2 + |\nabla u(\mathbf{x}, t)|^2 \leq C(1+|\mathbf{x}|)^{-2-\varepsilon}$ for some $C, \varepsilon > 0$, so we can interchange integration and differentiation according to the formula

$$\frac{d}{dt} \int_B u(\mathbf{x}, t) d\mathbf{x} = \int_B \partial_t u(\mathbf{x}, t) d\mathbf{x}$$

where B is any ball in \mathbf{R}^3 (see *Strauss*, p. 420, for the result for one spatial variable, but the same rule applies in \mathbf{R}^3). Show that if

$$\int_{\mathbf{R}^3} |u(\mathbf{x}, t_0)|^2 d\mathbf{x} = 1$$

for some $t_0 \in \mathbf{R}$, then

$$\int_{\mathbf{R}^3} |u(\mathbf{x}, t)|^2 d\mathbf{x} = 1$$

for all $t \in \mathbf{R}$.